



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 38/16, 38/18, 38/30, 38/17, 38/39 //</b> <b>(A61K 38/16, 39:395, 38:17, 31:70)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/55359</b> <b>(43) International Publication Date:</b> 4 November 1999 (04.11.99)
<b>(21) International Application Number:</b> PCT/US99/08303 <b>(22) International Filing Date:</b> 15 April 1999 (15.04.99) <b>(30) Priority Data:</b> 60/083,472                      29 April 1998 (29.04.98)                      US <b>(71) Applicant:</b> ALLERGAN SALES, INC. [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US). <b>(72) Inventors:</b> DOLLY, J., Oliver; 7 Buckland Road, Cheam, Surrey SM2 7LP (GB). AOKI, Kei, Roger; 2 Ginger Lily Court, Coto de Caza, CA 92679 (US). DE PAIVA, Anton; 62 Ravensmede Way, London W4 1TF (GB). <b>(74) Agents:</b> FISHER, Carlos, A. et al.; Allergan Sales, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> COMPOSITIONS AND METHODS FOR EXTENDING THE ACTION OF CLOSTRIDIAL NEUROTOXIN		
<b>(57) Abstract</b>  Methods and compositions for modulating neurite outgrowth in damaged neural endplates. Also disclosed are methods for extending the period during which tissue treated with Clostridial toxin is paralysed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## COMPOSITIONS AND METHODS FOR EXTENDING THE ACTION OF CLOSTRIDIAL NEUROTOXIN

This application claims priority under 35 U.S.C.  
5 §119(e)(1) to provisional application number 60/083,472,  
filed 4/29/98.

Field of the Invention

10 The present invention is directed towards methods and  
compositions for inhibiting neural sprouting in neurons that  
have been subjected to botulinum toxin. Also disclosed are  
methods and compositions for extending the period of time  
during which treatment of nerve cells with botulinum toxin  
15 is effective to prevent innervation of a cell or tissue,  
such as muscle cells or tissue. Such methods and  
compositions are effective in the treatment of spasms or  
muscular tetanus. Disclosed as well are methods and  
compositions for stimulating neural outgrowth.

20

Background of the invention

Neurotoxins, such as those obtained from *Clostridium*  
*botulinum* and *Clostridium tetanus*, are highly potent and  
specific poisons of neural cells. These Gram positive  
25 bacteria secrete two related but distinct toxins, each  
comprising two disulfide-linked amino acid chains: a light  
chain (L) of about 50 KDa and a heavy chain (H) of about 100  
KDa, which are wholly responsible for the symptoms of these  
diseases.

The tetanus and botulinum toxins are among the most lethal substances known to man, having a lethal dose in humans of between 0.1 ng and 1 ng per kilogram of body weight. Tonello et al., *Adv. Exp. Med. & Biol.* 389:251-260 (1996). Both toxins function by inhibiting neurotransmitter release in affected neurons. The tetanus neurotoxin (TeNT) acts mainly in the central nervous system, while botulinum neurotoxin (BoNT) acts at the neuromuscular junction by inhibiting acetylcholine release from the axon of the affected neuron into the synapse, resulting in a localized flaccid paralysis. The effect of intoxication on the affected neuron is long-lasting and has been thought to be irreversible.

The tetanus neurotoxin (TeNT) is known to exist in one immunologically distinct type; the botulinum neurotoxins (BoNT) are known to occur in seven different immunogenic types, termed BoNT/A through BoNT/G. While all of these types are produced by isolates of *C. botulinum*, two other species, *C. baratii* and *C. butyricum* also produce toxins similar to /F and /E, respectively. See e.g., Coffield et al., *The Site and Mechanism of Action of Botulinum Neurotoxin in Therapy with Botulinum Toxin* 3-13 (Jankovic J. & Hallett M. eds. 1994), the disclosure of which is incorporated herein by reference.

Regardless of type, the molecular mechanism of intoxication appears to be similar. In the first step of the process, the toxin binds to the presynaptic membrane of the target neuron through a specific interaction between the

heavy chain and a cell surface receptor; the receptor is thought to be different for each type of botulinum toxin and for TeNT. The carboxy terminus of the heavy chain appears to be important for targeting of the toxin to the cell

5 surface.

In the second step, the toxin crosses the plasma membrane of the poisoned cell. The toxin is first engulfed by the cell through receptor-mediated endocytosis, and an endosome containing the toxin is formed. The toxin then  
10 escapes the endosome into the cytoplasm of the cell. This last step is thought to be mediated by the amino terminus of the heavy chain, which triggers a conformational change of the toxin in response to a pH of about 5.5 or lower.

Endosomes are known to possess a proton pump which decreases  
15 intra-endosomal pH. The conformational shift exposes hydrophobic residues in the toxin, which permits the toxin to embed itself in the endosomal membrane. The toxin then translocates through the endosomal membrane into the cytosol.

20 The last step of the mechanism of botulinum toxin activity appears to involve reduction of the disulfide bond joining the heavy and light chain. The entire toxic activity of botulinum and tetanus toxins is contained in the light chain of the holotoxin; the light chain is a zinc  
25 (Zn++) endopeptidase which selectively cleaves proteins essential for recognition and docking of neurotransmitter-containing vesicles with the cytoplasmic surface of the plasma membrane, and fusion of the vesicles with the plasma

membrane. TxNT, BoNT/B BoNT/D, BoNT/F, and BoNT/G cause degradation of synaptobrevin 2 (also called vesicle-associated membrane protein (VAMP)), a synaptosomal membrane protein. Most of the VAMP present at the cytosolic surface of the synaptic vesicle is removed as a result of any one of these cleavage events. Each toxin specifically cleaves a different bond.

BoNT/A and /E selectively cleave the plasma membrane-associated protein SNAP-25; this protein is bound to and present on the cytosolic surface of the plasma membrane. BoNT/C cleaves syntaxin, an integral protein having most of its mass exposed to the cytosol. Syntaxin interacts with the calcium channels at presynaptic terminal active zones. See Tonello et al., *Tetanus and Botulism Neurotoxins in Intracellular Protein Catabolism* 251-260 (Suzuki K & Bond J. eds. 1996), the disclosure of which is incorporated by reference as part of this specification. BoNT/C<sub>1</sub> also cleaves SNAP-25 at a peptide bond next to that cleaved by BoNT/A.

Both TeNT and BoNT are taken up at the neuromuscular junction. BoNT remains within peripheral neurons, and blocks release of the neurotransmitter acetylcholine from these cells. Through its receptor, TeNT enters vesicles that move in a retrograde manner along the axon to the soma, and is discharged into the intersynaptic space between motor neurons and the inhibitory neurons of the spinal cord. At this point, TeNT binds receptors of the inhibitory neurons, is again internalized, and the light chain enters the

cytosol to block the release of the inhibitory neurotransmitters 4-aminobutyric acid (GABA) and glycine from these cells. Id.

Because of its specifically localized effects, dilute preparations of BoNT have been used since 1981 as therapeutic agents in the treatment of patients having various spastic conditions, including strabismus (misalignment of the eye), bophlarospasm (involuntary eyelid closure) and hemifacial spasm. See e.g., Borrodic et al., *Pharmacology and Histology Botulinum Toxin in Therapy with Botulinum Toxin 3-13* (Jankovic J. & Hallett M. eds. 1994), incorporated by reference herein. Of the seven toxin types, BoNT/A is the most potent of the BoNTs, and the most well characterized. Intramuscular injection of spastic tissue with dilute preparations of BoNT/A has been also used effectively to treat spasticity due to brain injury, spinal cord injury, stroke, multiple sclerosis and cerebral palsy.

The extent of paralysis depends on both the dose and dose volume delivered to the target site. Typically, the neurotoxin is administered in a preparation that also contains several non-toxic proteins as well, including hemagglutins and associated glycoproteins that assist in maximizing its stability and presentation to the target motor neuron.

Typically there is a 24 to 72 hour delay between the administration of the toxin and the onset of the clinical effect. Exposure to the toxin causes denervation atrophy. See e.g., Dutton J., *Acute and Chronic Effects of Botulinum*

*Toxin in the Management of Blepharospasm, in Therapy with Botulinum Toxin* at 199, incorporated by reference herein. Although the therapeutic application of BoNT has been extraordinarily effective, what side effects have been  
5 observed are mainly immediate effects associated with the treatment event itself. Peripheral effects causing weakness in adjacent muscles may sometimes occur; these effects usually do not persist beyond 1 to two weeks. The specific manifestations of these effects upon adjacent muscle groups  
10 depend upon the particular indication being treated. For example, patients treated for blepharospasm sometimes experience ptosis, and swallowing problems may occur after injection of neck muscles for torticollis.

Other possible consequences of treatment include the  
15 potential for overdosage through miscalculation or differences in activity between different preparations of the toxin, generalized fatigue, and the potential for allergic reactions.

A feature of treatment with BoNT/A, and other  
20 clostridial neurotoxin types, is that the paralytic action is temporary with symptoms reappearing in patients within a few months after toxin injection. This characteristic has been thought to be associated with the observed sprouting of nascent, synaptically active processes at the neuromuscular  
25 junction (NMJ). The production of such sprouts following BoNT/A therapeutic treatment has appeared to contribute to the reinnervation of the treated tissue and therefore the need for repeated serial injections of the toxin.



The duration of paralytic action caused by clostridial neurotoxin and the extent of sprouting, appears to depend on the neurotoxin subtype studied and therefore appears to be related to the specific SNARE target cleaved by each neurotoxin. Thus, BoNT/A and BoNT/C<sub>1</sub>, which cleave the t-SNARE protein SNAP-25 within one amino acid of each other, cause a long lasting paralysis with a long average sprout length observed. BoNT/F, which cleaves the v-SNARE protein VAMP, causes a paralysis of shorter duration and sprouts of shorter average length. BoNT/E, which cleaves SNAP-25 at a different position than that of BoNT/A and BoNT/C<sub>1</sub> (thereby liberating a SNAP fragment of different size from the plasma membrane), has a short duration period of about 5 days, and virtually no neural sprouting is observed. Without wishing to be bound by theory, these observations suggest that neural sprouting and duration of paralysis are normally related events, and that the cleavage products of BoNT proteolytic digestion (i.e., the liberated fragment or the membrane-bound fragment) can directly or indirectly regulate, or be coregulated with, neural sprouting.

It would therefore be advantageous to design a method whereby the sprouting phenomenon may be uncoupled from duration of therapeutic effect, and thus delayed, blocked or attenuated so as to prolong the effects of injection of tissue with toxin. Although the experiments described herein utilize a BoNT/A preparation, it will be recognized by those of skill in the art that the methods shown herein will be suitable for employment using other clostridial

toxins, such as BoNT/B through /G, and TeNT, in which a sprouting pathway can be observed.

Additionally, it would be advantageous to provide herein compositions effective for the inhibition or  
5 prevention of the sprouting phenomenon. Such compositions and methods would lessen the need for patients to undergo repeated neurotoxin treatment.

#### Summary of the Invention

10

The present invention concerns methods for increasing the period of time between therapeutic treatments of neural tissue with a clostridial neurotoxin; thus the method provides a method of increasing the effectiveness of such  
15 treatments. A direct advantage of such methods is an increased therapeutic "life" and a concomitant lessening in the required frequency of treatment of the patient with neurotoxin. Reducing frequency of treatment would provide less opportunity for a patient to experience the side  
20 effects described above that are observed following treatment, but which tend to subside long before the effectiveness of the toxin in the target area has subsided.

Additionally, reduced frequency of treatment provides less opportunity for miscalculation of dosage amount and other  
25 treatment-specific risks.

Accordingly, an aspect of the invention concerns a composition comprising a first agent comprising a clostridial neurotoxin for use as a therapeutic agent and a

second agent able to extend the duration of therapeutic benefit of said first agent, wherein the second agent is effective to attenuate the production of nerve terminal sprouts following treatment of a neuromuscular junction with  
5 the clostridial neurotoxin.

In a particular embodiment of this aspect of the invention, the first and second agent may comprise a single entity which is provided the patient in a single treatment session. For example, the entity may comprise a single  
10 molecule, or a disulfide-linked multichain polypeptide. Additionally or alternatively, the entity may comprise one or more adsorbed or linked heterogroup, such as a small organic molecule or a nucleic acid linked thereto. The entity preferably comprises both the receptor binding and  
15 translocation activities of a clostridial heavy chain and an active portion of a clostridial toxin light chain. The light chain may also comprise an auxiliary enzymatic activity, such as a ribonuclease, which specifically cleaves a nucleic acid encoding an intraneuronal factor which is  
20 responsible for the expression, activation and/or secretion of neurotrophic factors or cell adhesion molecules. In a preferred embodiment, such an auxiliary activity is provided by a ribozyme. By ribozyme is meant a nucleic acid or nucleic acid analog having a sequence-specific nuclease  
25 activity; the construction and use of ribozymes are well known in the art; see e.g., Cech, T., *Science* 236:1532-1539(1987); Cech, T. R., *Curr. Opin. Struct. Biol.* 2:605-609 (1992); and Usman et al., *Nucleic Acids & Mol. Biol.* 10:243-

264 (1996), the disclosures of which are hereby incorporated by reference herein. By nucleic acid analog is meant a polymeric molecule able to form a sequence-specific hybrid with a target single-stranded nucleic acid; such analogs may  
5 contain modified nucleotides (or ribonucleotides) such as 3'-O methyl nucleotides, phosphorothioate modified nucleotides, methylphosphonate nucleotides, or nucleotide bases separated by a peptide-like bond.

Alternatively, a nucleic acid or nucleic acid analog  
10 comprised in the single entity referred to above may be an antisense agent able to selectively bind to a nucleic acid encoding an intraneuronal factor which is responsible for the expression, activation and/or secretion of neurotrophic factors or cell adhesion molecules. This antisense agent  
15 may further provide a double-stranded substrate for the action of an intracellular RNase H activity. Details concerning certain embodiments of these aspects of the invention are contained in e.g., Dolly et al., International Publication No. WO95/32738, entitled *Modification of*  
20 *Clostridial Toxins for Use as Transport Proteins* and Uherek et al., *J. Biol. Chem.* 273:8835-8841 (1998). These two references are incorporated by reference as part of the present application.

A nucleic acid moiety linked polypeptide portion of the  
25 entity may encode a protein or polypeptide having the ability to be expressed within a neuron and to directly or indirectly regulate the expression, activation and/or secretion of neurotrophic factors or cell adhesion

molecules.

In preferred aspects of the invention, the second agent is selected from the group consisting of agents able to compete with, down-regulate, or neutralize the effects of:  
5 IGFI, IGF II, a neurotrophic factor, leukemia inhibitory factor, a nerve cell adhesion molecule and neural agrin. In a more preferred aspect of the invention, the neurotrophic factor is selected from the group consisting of: ciliary neurotrophic factor, NT-3, NT-4, and brain-derived  
10 neurotrophic factor and/or the nerve cell adhesion molecule is selected from the group consisting of tenascin-C, ninjurin, neural cell adhesion molecule.

Applicants have surprisingly discovered that recovery of neural function following poisoning of nerve terminals  
15 with clostridial neurotoxin involves two distinct and apparently coordinated events. First, the poisoned endplate becomes synaptically inactive. Shortly thereafter the endplate elaborates thin nascent axon neural processes. These processes or "sprouts" are synaptically competent  
20 after about 14 days following treatment with clostridial neurotoxin. The sprouts continue growing, reaching a maximal length and level of complexity after about 42 days following treatment with neurotoxin. During this time, the endplate remains synaptically inactive.

25 Secondly, after about 42 days, the sprouts begin to regress, shortening in length and decreasing in complexity.

At the same time, the original endplate begins to become synaptically active, undergoing synaptic vesicle turnover.

The increase in such turnover reaches that of the unpoisoned endplate after about 91 days post-treatment, at approximately the same time that the sprouting phenomenon has completely regressed, and no sprouts can be observed.

5 These findings are reported in DePaiva et al., *Proc. Nat'l. Acad. Sci.* 96:3200-3205 (March 1999), which is hereby incorporated by reference herein.

These observations are diametrically opposed to the prevailing wisdom in the art, in which it has largely been

10 assumed that the original endplate is permanently inactivated upon treatment with clostridial toxin, and that all return of synaptic activity is due to extension of sprouts to compensate for the lack of a neurologically functional endplate. However, as indicated, the old

15 paradigm has been shown by the present Applicants to be in error, in that the poisoned endplate regains neurological function over time, while the axon sprouts regress so that after a given time period the nerve terminal appears essentially as it did prior to treatment. Therefore,

20 Applicants have discovered that, far from being a permanent feature, the axonal sprouts assume a temporary role in the rehabilitation of the poisoned endplate.

Although not wishing to be limited by theory, Applicants believe that these results indicate that the two

25 events outlined above are temporally coordinated, in that a blockage of the neural sprouting phenomenon would delay or block the recovery of the functional endplate. Such temporal coordination could be due to the secretion of one

or more factor by the damaged neural endplate (or the inactive muscle fiber) that has neurotrophic effects resulting in the formation of neural sprouts; these sprouts may then elaborate a factor (either the same or different  
5 from the first factor) which promotes continued sprouting. This factor may be produced during neurite sprouting in amounts sufficient for the reinnervation of the neurotoxin-damaged endplate even after treatment with clostridial toxin. Thus, treatment of cells with an agent able to block  
10 neural sprouting would also delay or otherwise attenuate the ability of the treated endplate to experience return of neurological function, and in fact may well block such return altogether.

As indicated, the signaling event indicating the  
15 initiation of the neural sprouting phenomenon appears to be mediated by a cytokine or other intercellular messenger. One such agent, agrin, appear to be an important player, if not the key molecule, in the formation of the neuromuscular junction in development, and in neuromuscular regeneration.  
20 See Ruegg M.A. and Bixby J. L., *Trends in Neurol. Sci.* 21:22-27 (1998), the disclosure of which is incorporated herein by reference. Agrin appears to be present in a number of isoforms, which result from alternative mRNA splicing. Soluble agrin isolated from synaptic basal lamina extracts  
25 (to which it binds following secretion) is able to induce the aggregation of acetylcholine receptors in the postsynaptic portions of muscle cells. Agrin present in motor neuron terminals (n-agrin) contains an insert,

relative to other agrin species, in a region termed the B/z. region; this insert is important in conferring the ability on n-agrin to aggregate acetylcholine receptors in postsynaptic tissue. Neural agrin is released by the motor-  
5 nerve terminal and is believed by Applicants to induce post-synaptic specialization and up-regulation of other factors, such as muscle-diffusible factors, involved in the neural sprouting response.

It is anticipated that methods which interfere with the  
10 sprouting phenomenon (such as by the preventing the action of agrin) would delay the return of innervation to cells which are controlled by neurons which have been therapeutically poisoned (i.e., by BONT or TeNT) or otherwise damaged. This is because, as indicated above, the  
15 rate of return of neural function appears to be partly dependent upon the presence of synaptically active sprouts.

Additionally, agents to be used in the inhibition of sprouting may very well also delay or prevent the recovery of neural activity of the endplate following poisoning.

20 Thus utilization of such methods would therefore be expected to extend the effective period of treatment of tissue with a clostridial toxin, by delaying the regeneration of active neuromuscular synapses, both through inhibition of sprouting and of recovery of the poisoned  
25 endplate.

#### Detailed Description of the Invention



This invention is drawn to methods and compositions for increasing the therapeutic effectiveness of treatment of tissue with clostridial neurotoxin. This increase in effectiveness is made possible by the surprising discovery  
5 that regeneration of neural tissue damaged by treatment with clostridial neurotoxin is a complex occurrence in which two coordinated events take place.

In the first of these events, the poisoned neuromuscular endplate becomes synaptically inactive,  
10 demonstrating no exocytosis of synaptic vesicles and thus no transport of intracellular acetylcholine. In four days after treatment, the endplate begins to form neural sprouts that are shown to release and regenerate synaptic vesicles.

These sprouts grow in length and complexity until  
15 approximately 42 days following treatment with the neurotoxin; at this point the neural sprouts begin to regress and shorten. At ninety-one days following treatment, the neural sprouts can no longer be seen.

In the second event, simultaneously with the beginning  
20 of regression of the neural sprouts, the synaptically inactive endplate begins to regain the ability to release acetylcholine and begin to recycle synaptic vesicles. This ability, which begins at relatively low levels, increases over the time period indicated above. At approximately 91  
25 days following treatment with clostridial neurotoxin the endplate is histologically and synaptically indistinguishable from the condition of the endplate before treatment with clostridial neurotoxin.

These findings indicate that one may therapeutically intervene at one of the major steps of the sprouting phenomenon to prevent or attenuate the neural sprouting as a method of extending the effective period during which tissue treated with the toxin remains paralyzed. In an initial step, the muscle cells surrounding the neural endplate respond, either sensing the inactive muscle or in response to a signal from the poisoned nerve terminal, by producing muscle-derived diffusable factors. The expression of a number of muscle-derived signaling factors appears to be upregulated by muscle inactivation; such factors include insulin-like growth factors (IGF-1 and IGF-2). A factor such as neural agrin is believed to be the initial signal directing the muscle cell to produce the IGF molecules. Reports have demonstrated that IGF I and IGF II effect neurite outgrowth in cultured BoNT/A treated dorsal root ganglia, and also are able to stimulate the initial sprouting response in paralyzed mouse gluteus muscle. See Caroni, P. and Schneider, C. *J. Neurosci.* 14:3378-3388 (1994) and Caroni, P., et al. *J. Cell Biol.* 125:893-902 (1994).

Thus, blocking the effects of such muscle derived diffusable factors that positively affect neurite outgrowth and sprouting may attenuate not only clostridial neurotoxin-induced sprouting, but may also delay the eventual recovery of neurotransmission at the poisoned nerve terminals. Such blocking may occur through the use of antibodies specific for the muscle derived diffusable factor in question, or

that are common to such muscle derived diffusable factors. .  
 Alternatively, there are naturally occurring binding  
 proteins, such as the IGF binding proteins IGF-BP 4 and IGF-  
 BP 5, which can bind to, and therefore block, the  
 5 neurotrophic effect of such diffusable factors.

IGF-BP 4 has an amino acid sequence (from the amino  
 terminus) of:

10 MLPLCLVAALLLAAGPGPSLGDEAIHCPPCSEKLRARPPVGCCELVREPGCGCCATCA  
 LGLGMPCGVYTPRCGSLRCYPPRGVEKPLHTLMHGQGVCMELAEIEAIQESLQPSDKDE  
 GDHPNNSFSPCSAHDRRLQKHFAKIRDRSTSGGKMKVNGAPREDARPVPOGSCQSELHR  
 ALERLAASQSRTHEDLYIIPINCDRNGNFHPKQCHPALDGQRGKCWCVDKRTGVKLPGG  
 LEPKGELDCHQLADSFRE

15 (SEQ ID NO: 1)

This is encoded within the nucleotide base sequence (from 5'  
 to 3') of:

20 GTGCCCTCCG CCGCTCGCCC GCGCGCCCGC GCTCCCCGCC TCGCGCCAGC  
 GCCCCGCGCC CGCGCCCCAG TCCTCGGGCG GTCATGCTGC CCCTCTGCCT  
 CGTGCCCGCC CTGCTGCTGG CCGCCGGGCG CCGGCCGAGC CTGGGCGACG  
 AAGCCATCCA CTGCCCAGCC TGCTCCGAGG AGAAGCTGGC GCGCTGCCGC  
 CCCCCGTGG GCTGCGAGGA GCTGGTGCGA GAGCCGGGCT GCGGCTGTTG  
 25 CGCCACTTGC GCCCTGGGCT TGGGGATGCC CTGCGGGGTG TACACCCCCC  
 GTTGCGGCTC GGGCCTGCGC TGCTACCCGC CCCGAGGGGT GGAGAAGCCC  
 CTGCACACAC TGATGCACGG GCAAGGCGTG TGCATGGAGC TGGCGGAGAT  
 CGAGGCCATC CAGGAAAGCC TGCAGCCCTC TGACAAGGAC GAGGGTGACC  
 ACCCCAACAA CAGCTTCAGC CCCTGTAGCG CCCATGACCG CAGGTGCCTG  
 30 CAGAAGCACT TCGCCAAAAT TCGAGACCGG AGCACCAGTG GGGGCAAGAT  
 GAAGGTCAAT GGGGCGCCCC GGGAGGATGC CCGGCCTGTG CCCCAGGGCT  
 CCTGCCAGAG CGAGCTGCAC CCGGCGCTGG AGCGGCTGGC CGCTTCACAG  
 AGCCGCACCC ACGAGGACCT CTACATCATC CCCATCCCCA ACTGCGACCG  
 CAACGGCAAC TTCCACCCCA AGCAGTGTCA CCCAGCTCTG GATGGGCAGC  
 35 GTGGCAAGTG CTGGTGTGTG GACCGGAAGA CCGGGGTGAA GCTTCCGGGG  
 GGCCTGGAGC CAAAGGGGGA GCTGGACTGC CACCAGCTGG CTGACAGCTT  
 TCGAGAGTGA GGCCTGCCAG CAGGCCAGGG ACTCAGCGTC CCCTGCTACT  
 CCTGTGCTCT GGAGGCTGCA GAGCTGACCC AGAGTGGAGT CTGAGTCTGA

```

GTCCTGTCTC TGCCTGCGGC CCAGAAGTTT CCCTCAAATG CGCGTGTGCA
CGTGTGCGTG TGCCTGCGTG TGTGTGTGTT TGTGAGCATG GGTGTGCCCT
TGGGGTAAGC CAGAGCCTGG GGTGTTCTCT TTGGTGTTAC ACAGCCCAAG
AGGACTGAGA CTGGCACTTA GCCCAAGAGG TCTGAGCCCT GGTGTGTTTC
5 CAGATCGATC CTGGATTAC TCACTCACTC ATTCCTTCAC TCATCCAGCC
ACCTAAAAAC ATTTACTGAC CATGTACTAC GTGCCAGCTC TAGTTTTTCAG
CCTTGGGAGG TTTTATTCTG ACTTCCTCTG ATTTTGGCAT GTGGAGACAC
TCCTATAAGG AGAGTTCAAG CCTGTGGGAG TAGAAAAATC TCATTCCCAG
AGTCAGAGGA GAAGAGACAT GTACCTTGAC CATCGTCCTT CCTCTCAAGC
10 TAGCCAGAGG GTGGGAGCCT AAGGAAGCGT GGGGTAGCAG ATGGAGTAAT
GGTCACGAGG TCCAGACCCA CTCCCAAAGC TCAGACTTGC CAGGCTCCCT
TTCTCTTCTT CCCCAGGTCC TTCCTTTAGG TCTGGTTGTT GCACCATCTG
CTTGTTGGC TGGCAGCTGA GAGCCCTGCT GTGGGAGAGC GAAGGGGGTC
AAAGGAAGAC TTGAAGCACA GAGGGCTAGG GAGGTGGGGT ACATTTCTCT
15 GAGCAGTCAG GGTGGGAAGA AAGAATGCAA GAGTGGACTG AATGTGCCTA
ATGGAGAAGA CCCACGTGCT AGGGGATGAG GGGCTTCCTG GGTCCCTGTT
CCTACCCCAT TTGTGGTCAC AGCCATGAAG TCACCGGGAT GAACCTATCC
TTCCAGTGGC TCGCTCCCTG TAGCTCTGCC TCCCTCTCCA TATCTCCTTC
CCCTACACCT CCCTCCCCAC ACCTCCCTAC TCCCCTGGGC ATCTTCTGGC
20 TTGACTGGAT GGAAGGAGAC TTAGGAACCT ACCAGTTGGC CATGATGTCT TTTCT

```

(SEQ ID NO: 3)

IGFBP 5 has an amino acid sequence (from the amino  
 25 terminus) of:

```

MVLLTAVLLLLLAAYAGPAQSLGSFVHCEPCDEKALSMCPPSPLGCELVKEPGCGCCMTCA
LAEGQSCGVYTERCAQGLRCLPRQDEEKPLHALLHGRGVCLNEKSYREQVKIERDSREHE
EPTTSEMAEETYSPIFRPKHTRISELKAEAVKKDRRKKLTQSKFVGGAENTAHPRIISA
30 PEMRQESEQGPCRRHMEASLQELKASPRMVPRVYLPNCDRKGIFYKRKQCKPSRGRKGI
CWCVDKYGMKLPMEYVDGDFQCHTFDSSNVE

```

(SEQ ID No. 2)

35 and is encoded within a nucleotide base sequence (from 5' to  
 3') of:

```

GGGGAAGA GCTAGGAAAG AGCTGCAAAG CAGTGTGGGC TTTTTCCTT
TTTTTGCTCCT TTTCATTAC CCCTCCTCCG TTTTACCCT TCTCCGGACT
40 TCGCGTAGAA CCTGCGAATT TCGAAGAGGA GGTGGCAAAG TGGGAGAAAA

```

```

GAGGTGTTAG GGT TTGGGGT TTTT TGT TTT TTGTTT TTTT TTAATTT
CTTGATTTCA ACATTTTCTC CCACCCTCTC GGCTGCAGCC AACGCCTCTT
ACCTGTTCTG CGGCGCCGCG CACCGCTGGC AGCTGAGGGT TAGAAAGCGG
GGTGTATTTT AGATTTTAAAG CAAAAATTTT AAAGATAAAT CCATTTTCTT
5 CTCCCACCCC CAACGCCATC TCCACTGCAT CCGATCTCAT TATTTCCGGT
GTTGCTTGGG GGTGAACAAT TTTGTGGCTT TTTT TCCCCT ATAATTCTGA
CCCGCTCAGG CTTGAGGGT TCTCCGGCCT CCGCTCACTG CGTGACCTG
GCGCTGCCCT GCTTCCCCCA ACCTGTTGCA AGGCTTTAAT TCTTGCAACT
GGGACCTGCT CGCAGGCACC CCAGCCCTCC ACCTCTCTCT ACATTTTTCG
10 AAGTGTCTGG GGGAGGGCAC CTGCTCTACC TGCCAGAAAT TTTAAACAA
AAACAAAAAC AAAAAAATCT CCGGGGGCCC TCTTGGCCCC TTTATCCCTG
CACTCTCGCT CTCCTGCCCC ACCCCGAGGT AAAGGGGGCG ACTAAGAGAA
GATGGTGTTG CTCACCGCGG TCCTCTGCT GCTGGCCGCC TATGCGGGGC
CGGCCAGAG CCTGGGCTCC TTCGTGCACT GCGAGCCCTG CGACGAGAAA
15 GCCCTCTCCA TGTGCCCCC CAGCCCCCTG GGCTGCGAGC TGGTCAAGGA
GCCGGGCTGC GGCTGCTGCA TGACCTGCGC CCTGGCCGAG GGGCAGTCGT
GCGGCGTCTA CACCGAGCGC TGCGCCAGG GGCTGCGCTG CCTCCCCCGG
CAGGACGAGG AGAAGCCGCT GCACGCCCTG CTGCACGGCC GCGGGGTTTG
CCTCAACGAA AAGAGCTACC GCGAGCAAGT CAAGATCGAG AGAGACTCCC
20 GTGAGCACGA GGAGCCCACC ACCTCTGAGA TGGCCGAGGA GACCTACTCC
CCCAAGATCT TCCGGCCCAA ACACACCCGC ATCTCCGAGC TGAAGGCTGA
AGCAGTGAAG AAGGACCGCA GAAAGAAGCT GACCCAGTCC AAGTTTGTG
GGGGAGCCGA GAACACTGCC CACCCCCGGA TCATCTCTGC ACCTGAGATG
AGACAGGAGT CTGAGCAGGG CCCCTGCCGC AGACACATGG AGGCTTCCCT
25 GCAGGAGCTC AAAGCCAGCC CACGCATGGT GCCCCGTGCT GTGTACCTGC
CCAATTGTGA CCGCAAAGGA TTCTACAAGA GAAAGCAGTG CAAACCTTCC
CGTGGCCGCA AGCGTGGCAT CTGCTGGTGC GTGGACAAGT ACGGGATGAA
GCTGCCAGGC ATGGAGTACG TTGACGGGGA CTTTCAGTGC CACACCTTCG
ACAGCAGCAA CGTTGAGTGA TCGTCCCCC CCAACCTTT CCCTCACCCC
30 CTCCCACCCC CAGCCCCGAC TCCAGCCAGC GCCTCCCTCC ACCCCAGGAC
GCCACTCATT TCATCTCATT TAAGGGAAAA ATATATATCT ATCTATTGA
GGAAAAAAAA AAAAAAAAAA AA

```

(SEQ ID NO: 4)

35

These binding proteins can be made synthetically or cloned and produced for therapeutic purposes, while a cell line producing a desired monoclonal antibody can be maintained for relatively large-scale antibody production.

40

Cloning and general antibody methodologies are commonplace

in the art; such methodologies are disclosed within Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd ed. Cold Spring Harbor Laboratory Press 1989), the disclosure of which is hereby incorporated by reference as part of this disclosure.

Rather than using competitive methods, another aspect of the invention involves the use of a cholinergic special transporter to insert a gene which produces inactive receptors for one or more factor involved in promotion of neural sprouting. Such receptors would maintain high specific binding constants to their ligands, but the biological activity of the receptors would be abrogated; such receptors could be easily be generated and screened through the introduction of mutations in the nucleotide sequence encoding the protein, and assaying the mutants for binding strength and biological activity. Alternatively, the neurotrophic activities of factors produced by the neural endplate or nascent sprouts may be inhibited through intracellular targeting and delivery of a competitive inhibitor, ribozyme, transcriptional suppresser or other agent specifically able to block or attenuate such activities, as described above. Such intracellular targeting is disclosed in references such as, e.g., Dolly et al., International Patent Publication No. WO95/32738, previously incorporated by reference herein.

The second point at which intervention in the sprouting phenomenon may be made is during the stage of axonal outgrowth and arbor development. At this stage such

outgrowth has already been initiated, but auxiliary factors appear to be necessary in order to maintain axonal growth. A number of factors are known to affect the rates of outgrowth, and may also effect the survivability of many types of neurons. Such factors include ciliary neurotrophic factor (CNTF); neurotrophins, including NT-3, NT-4, and brain-derived neurotrophic factor (BDNF); and leukemia inhibitory factor (LIF). Not only are these factors important in establishing an initial rate of axonal outgrowth, but they appear to either directly or indirectly stimulate their own production -- therefore blocking the initial outgrowth of sprouts may be essential in preventing further propagation through the continued expression of such factors.

15       The same methods as disclosed above can be used to prevent one or more of the agents listed above to manifest its activity. As would be expected by those of skill in the art in light of the present disclosure, such agents, destroy or bind to, and therefore attenuate the activity of these factors or their block their ability to bind to their receptors, or inactive forms of their receptors, would be useful when used in conjunction with a clostridial neurotoxin to prevent or reduce the rate of sprouting of neurites from the poisoned or damaged endplate.

25       The third stage at which the sprouting phenomenon may be attenuated or inhibited concerns the binding of axons to the extracellular matrix. Axons are guided to cellular processes containing the appropriate neurotransmitter

receptors by binding to components of the extracellular matrix. Such binding involves a variety of cell-borne or matrix associated adhesion molecules. Tenascin-C is an extracellular matrix component derived from Schwann cells that appears to bind neural processes. Ninjurin is a cell surface adhesion molecule that is up-regulated following peripheral nerve injury and thought to be involved in axonal guidance. See Araki, T., et al., *J. Biol. Chem.* 272:21373-21380 (1997), incorporated by reference herein. Similarly, neural-cell adhesion molecule (N-CAM), is an adhesion molecule which is thought to be involved in binding of neural sprouts to the extracellular matrix. The same techniques indicated above may be employed to prevent the expression or inhibit the activity of these molecules when used as part of a clostridial neurotoxin therapy.

Thus, in one aspect the present invention is drawn to a method for extending the effective period during which tissue treated with clostridial toxin is paralyzed, comprising: Contacting said tissue with a composition comprising an agent able to prevent the neuroregenerative activity of a polypeptide selected from the group consisting of IGF-1, IGF-2, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin.

In one preferred embodiment, the agent comprises a polypeptide able to bind to IGF-1 and/or IGF-2 in a manner that prevents an IGF molecule from binding to or activating



a cell surface receptor involved in the initiation of neural sprouting. In a most preferred embodiment the polypeptide comprises at least a portion of a amino acid sequence selected from the group consisting of: IGFBP-4 (SEQ ID NO: 1) or IGFBP5 (SEQ ID NO: 2). Preferably said portion comprises at least 10 contiguous amino acids of said sequence; more preferably said portion comprises at least 20 contiguous nucleotides of said sequence. Most preferably, the portion comprises an amino acid sequence selected from the group consisting of the entire amino acid sequence of IGFBP-4 or IGFBP-5.

Treatment of cells with such a composition may be accomplished either before or simultaneously with treatment with clostridial toxin. Preferably, the clostridial toxin is a botulinum toxin. Even more preferably, the botulinum toxin comprises BoNT/A. In other embodiments, the clostridial toxin is TeNT. Agent which are able to bind to any of these factors in a manner that inhibits their neurotrophic activity, or which bind to the receptors for such factors, would, in light of the present application, be expected to function as agents for extending the effective period between treatments of tissue with a neurotoxin.

Another embodiment comprises a cholinergic specific transporter joined to a gene encoding a gene which produces an inactive receptor for one or more factor involved in promotion of neural sprouting when delivered to a neural cell in vivo. Preferably the receptors maintain high specific binding constants to said factor(s) and the

biological activity of the receptor is reduced or absent.  
In a particularly preferred embodiment the transporter  
comprises some or all of a clostridial neurotoxin heavy  
chain, although other transporters such as the diphtheria  
5 toxin transporter may be effective in this regard as well.

In another embodiment the invention comprises a  
cholinergic specific transporter that is covalently or non-  
covalently joined to a nucleic acid which comprises a  
ribozyme or antisense nucleic acid able to specifically  
10 destroy the nucleic acids encoding neurotrophic agents or  
their receptors. Said joining may be made through methods  
including, but not limited to, covalent bonding or  
electrostatic forces.

In another embodiment, the present invention is drawn  
15 to the methods for stimulating the outgrowth of neural  
sprouts from damaged neural tissue. Such methods could be  
effective ways of increasing the rate at which reinnervation  
occurs after a neural injury. These methods comprise:  
Contacting said tissue with a composition comprising a  
20 polypeptide which comprises a neurotrophically active domain  
derived from an agent selected from a group consisting of  
IGF-1, IGF-2, ciliary neurotrophic factor, NT-3, NT-4, brain  
derived neurotrophic factor, leukemia inhibitory factor,  
tenascin-C, ninjurin, neural cell adhesion molecule, and  
25 neural agrin. Such damage may be a result of neurotoxin  
poisoning or due to a traumatic event, including but not  
limited to nerve or spinal cord crush injuries, traumatic  
brain injuries, glaucoma-induced damage to the retina and/or

optic nerve, or surgical trauma or injury.

The following examples illustrate various embodiments of the present invention, and are not intended to limit the scope of the invention, which is solely defined by the  
5 claims which conclude this specification.

#### Example 1

Blepharospasm is a medical condition characterized by  
10 uncontrolled eyelid movement. In its early stages, the condition is characterized by excessive blinking or fluttering of the eyelids. The condition is generally a progressive one, in which excessive blinking is replaced in the later stages with spasms of eye closure that interfere  
15 with visual function. The spasms become more frequent and severe, and involve the preseptal, pretarsal, and orbicularis oculi muscle. The condition often results in functional blindness relatively quickly (in a matter of two to three years) after the symptoms are first encountered.

20 A patient suffering from moderate idiopathic blepharospasm is treated with injections of BoNT/A toxin preparation containing non-toxic proteins and hemagglutins in sterile saline. Alternatively, the same toxin preparation without hemagglutins may be used. The  
25 injections are generally in the volume of 100  $\mu$ l; and each injection contains 1.25 to 2.5 units of the toxin preparation. The injections are made into the pretarsal orbicularis oculi of the upper lid laterally and medially

and in the lower lid laterally and medially. Additionally,  
2.5 unit injections (100  $\mu$ l each) are made lateral to the  
lateral canthus and into the brow medially. Total amount of  
BoNT/A toxin injected is roughly 6.25 to 12.5 units per  
5 eye. The Bo/A toxin is provided in a sterile, preservative-  
free saline, and the same solution is used to dilute the  
BoNT/A toxin if the master preparation of it is too  
concentrated.

Following injection the treated muscles are  
10 sufficiently paralyzed due to this treatment to alleviate  
the major symptoms of blepharospasm. Some mild concomitant  
weakness in the surrounding muscle tissue is observed; these  
side effects are mild and tolerated well by the patient.  
The effect of this treatment lasts approximately 8 weeks,  
15 and must be repeated at the end of this time to maintain the  
beneficial effects.

Example 2

A patient with blepharospasm is pre-treated with BoNT/A toxin as indicated in Example 1 with the following  
5 difference. Prior to injection with BoNT/A toxin preparation, the patient is given a ten-fold excess of IGFBP-4, having an amino acid sequence of SEQ ID NO: 1. The binding protein preparation is dissolved in sterile, preservative-free saline. Each injection is in the same  
10 area as the toxin injections that follow the pre-treatment; the volume of each injection is 100 µl. The BoNT/A toxin preparation is injected ten minutes after the injection of the IGFBT 4 injection.

The patient's therapeutic response to the BoNT/A toxin  
15 is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended to 12 weeks or more, during which time no further injection need be made.

20

Example 3

A patient with blepharospasm is pre-treated with BoNT/A toxin as indicated in Example 1 with the following  
25 difference. The BoNT/A toxin has been modified to have joined thereto a nucleic acid comprising a ribozyme specifically targeted to enzymatically destroy neural agrin mRNA. No supplemental injections are made.

The patient's therapeutic response to the BoNT/A toxin is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended to 12 weeks or more, during which time no further injection  
5 need be made.

#### Example 4

A patient with blepharospasm is pre-treated with BoNT/A  
10 toxin as indicated in Example 1 with the following difference. The BoNT/A toxin has been modified to have joined thereto a nucleic acid encoding an inactive neurotrophin receptor which retains the ability to bind its target neurotrophin. No supplemental injections are made.

15 The patient's therapeutic response to the BoNT/A toxin is similar to that seen in Example 1. The duration of the benefit provided by the BoNT/A toxin treatment is extended beyond that seen with BoNT/A alone, during which time no further injection need be made.

20 It will be understood that, while reference is made to BoNT/A in the Examples above, any other of the species of botulinum toxins (e.g., BoNT/B through G) could be substituted therefor, with appropriate adjustments possibly necessary due to differences in specific activity of the  
25 toxin. Additionally, the light chain segment could be derived from any clostridial neurotoxin (or other neurotoxin) , with the heavy chain retaining the motor

neuron receptor binding and exo-vesicular transport activities retained from the BoNT heavy chain.

These Examples are not intended to exhaust the embodiments of the present invention, and the invention is  
5 not to be seen as limited thereby. Further embodiments will be disclosed within the claims that conclude this specification.

CLAIMS

What is claimed is:

- 5 1) A method for extending the effective period during which tissue treated with a clostridial toxin is paralyzed comprising: contacting said tissue with a composition comprising an agent able to prevent the neuroregenerative activity of a polypeptide selected from the group consisting  
10 of: IGF I, IGF II, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin.
- 15 2) The method of claim 1 wherein said contacting step occurs at the same time as said tissue is treated with said clostridial toxin.
- 3) The method of claim 1 wherein said contacting step  
20 occurs prior to treatment of said tissue with said clostridial toxin.
- 4) The method of claim 1 wherein said clostridial toxin comprises BoNT.
- 25 5) The method of claim 1 wherein said clostridial comprises BoNT/A.



- 6) The method of claim 1 wherein said agent is selected from the group consisting of:
- a) an antibody able to selectively bind said polypeptide,
  - b) a competitive inhibitor of said polypeptide,
  - 5 c) a compound able to selectively prevent the expression of a gene encoding said polypeptide,
  - d) a binding protein other than an antibody, and
  - e) a ribozyme,
  - f) a nucleic acid encoding an inactive growth factor
- 10 receptor able to bind said growth factor.
- 7) The method of claim 6 wherein said agent is an antibody able to selectively bind said polypeptide.
- 15 8) The method of claim 6 wherein said agent is a competitive inhibitor of said polypeptide.
- 9) The method of claim 6 wherein said agent is a compound able to prevent the expression of a gene encoding said
- 20 polypeptide.
- 10) The method of claim 6 wherein said agent is a binding protein other than an antibody.
- 25 11) The method of claim 9 wherein said polypeptide is selected from the group consisting of IGF I and IGF II, and said binding protein is selected from the group consisting of IGF-BP4 and IGF-BP5.

- 12) A method for stimulating the outgrowth of neural sprouts from damaged neural tissue comprising: contacting said tissue with a composition comprising a polypeptide  
5 which comprises a neurotrophically active domain derived from an agent selected from the group consisting of IGF I, IGF II, ciliary neurotrophic factor, NT-3, NT-4, brain-derived neurotrophic factor, leukemia inhibitory factor, tenascin-C, ninjurin, neural cell adhesion molecule, and neural agrin.
- 10
- 13) The method of claim 11 wherein said agent comprises IGF I .
- 14) The method of claim 11 wherein said agent comprises IGF  
15 II.
- 15) The method of claim 11 wherein said agent comprises NT-3.
- 20 16) The method of claim 11 wherein said agent comprises ciliary neurotrophic factor.
- 17) The method of claim 11 wherein said agent comprises NT-3.
- 25
- 18) The method of claim 11 wherein said agent comprises NT-4.

19) The method of claim 11 wherein said agent comprises brain-derived neurotrophic factor.

20) The method of claim 11 wherein said agent comprises  
5 leukemia inhibitory factor.

21) The method of claim 11 wherein said agent comprises tenascin-C.

10 22) The method of claim 11 wherein said agent comprises ninjurin.

23) The method of claim 11 wherein said agent comprises neural-cell adhesion molecule.

15

24) The method of claim 11 wherein said agent comprises neural agrin.

## SEQUENCE LISTING

<110> Dolly, Oliver J.  
Aoki, Kei Roger  
De Paiva, Anton

<120> Compositions and Methods for Modulating  
Neural Sprouting

<130> 17259

<150> 60/083,472

<151> 1998-04-29

<160> 4

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 258

<212> PRT

<213> Homo sapiens

<400> 1

```

Met Leu Pro Leu Cys Leu Val Ala Ala Leu Leu Leu Ala Ala Gly Pro
 1           5           10           15
Gly Pro Ser Leu Gly Asp Glu Ala Ile His Cys Pro Pro Cys Ser Glu
          20           25           30
Glu Lys Leu Ala Arg Cys Arg Pro Pro Val Gly Cys Glu Glu Leu Val
          35           40           45
Arg Glu Pro Gly Cys Gly Cys Cys Ala Thr Cys Ala Leu Gly Leu Gly
          50           55           60
Met Pro Cys Gly Val Tyr Thr Pro Arg Cys Gly Ser Gly Leu Arg Cys
65           70           75           80
Tyr Pro Pro Arg Gly Val Glu Lys Pro Leu His Thr Leu Met His Gly
          85           90           95
Gln Gly Val Cys Met Glu Leu Ala Glu Ile Glu Ala Ile Gln Glu Ser
          100          105          110
Leu Gln Pro Ser Asp Lys Asp Glu Gly Asp His Pro Asn Asn Ser Phe
          115          120          125
Ser Pro Cys Ser Ala His Asp Arg Arg Cys Leu Gln Lys His Phe Ala
          130          135          140
Lys Ile Arg Asp Arg Ser Thr Ser Gly Gly Lys Met Lys Val Asn Gly
145          150          155          160
Ala Pro Arg Glu Asp Ala Arg Pro Val Pro Gln Gly Ser Cys Gln Ser
          165          170          175
Glu Leu His Arg Ala Leu Glu Arg Leu Ala Ala Ser Gln Ser Arg Thr
          180          185          190
His Glu Asp Leu Tyr Ile Ile Pro Ile Pro Asn Cys Asp Arg Asn Gly
          195          200          205
Asn Phe His Pro Lys Gln Cys His Pro Ala Leu Asp Gly Gln Arg Gly
          210          215          220
Lys Cys Trp Cys Val Asp Arg Lys Thr Gly Val Lys Leu Pro Gly Gly
225          230          235          240
Leu Glu Pro Lys Gly Glu Leu Asp Cys His Gln Leu Ala Asp Ser Phe
          245          250          255

```

Arg Glu

<210> 2  
 <211> 272  
 <212> PRT  
 <213> Homo Sapiens

<400> 2  
 Met Val Leu Leu Thr Ala Val Leu Leu Leu Leu Ala Ala Tyr Ala Gly  
 1 5 10 15  
 Pro Ala Gln Ser Leu Gly Ser Phe Val His Cys Glu Pro Cys Asp Glu  
 20 25 30  
 Lys Ala Leu Ser Met Cys Pro Pro Ser Pro Leu Gly Cys Glu Leu Val  
 35 40 45  
 Lys Glu Pro Gly Cys Gly Cys Cys Met Thr Cys Ala Leu Ala Glu Gly  
 50 55 60  
 Gln Ser Cys Gly Val Tyr Thr Glu Arg Cys Ala Gln Gly Leu Arg Cys  
 65 70 75 80  
 Leu Pro Arg Gln Asp Glu Glu Lys Pro Leu His Ala Leu Leu His Gly  
 85 90 95  
 Arg Gly Val Cys Leu Asn Glu Lys Ser Tyr Arg Glu Gln Val Lys Ile  
 100 105 110  
 Glu Arg Asp Ser Arg Glu His Glu Glu Pro Thr Thr Ser Glu Met Ala  
 115 120 125  
 Glu Glu Thr Tyr Ser Pro Lys Ile Phe Arg Pro Lys His Thr Arg Ile  
 130 135 140  
 Ser Glu Leu Lys Ala Glu Ala Val Lys Lys Asp Arg Arg Lys Lys Leu  
 145 150 155 160  
 Thr Gln Ser Lys Phe Val Gly Gly Ala Glu Asn Thr Ala His Pro Arg  
 165 170 175  
 Ile Ile Ser Ala Pro Glu Met Arg Gln Glu Ser Glu Gln Gly Pro Cys  
 180 185 190  
 Arg Arg His Met Glu Ala Ser Leu Gln Glu Leu Lys Ala Ser Pro Arg  
 195 200 205  
 Met Val Pro Arg Ala Val Tyr Leu Pro Asn Cys Asp Arg Lys Gly Phe  
 210 215 220  
 Tyr Lys Arg Lys Gln Cys Lys Pro Ser Arg Gly Arg Lys Arg Gly Ile  
 225 230 235 240  
 Cys Trp Cys Val Asp Lys Tyr Gly Met Lys Leu Pro Gly Met Glu Tyr  
 245 250 255  
 Val Asp Gly Asp Phe Gln Cys His Thr Phe Asp Ser Ser Asn Val Glu  
 260 265 270

<210> 3  
 <211> 1955  
 <212> DNA  
 <213> Homo Sapiens

<400> 3  
 gtgccctcgc ccgctcgccc gcgcgcccgc gctccccgcc tgcgcccagc gccccgcgcc  
 60  
 cgcgccccag tctcggggcg gtcattgtgc cctctgcct cgtggccgcc ctgctgtctg  
 120  
 ccgcccggggc cgggcccagc ctgggcgacg aagccatcca ctgcccgcgc tgctccgagg  
 180

agaagctggc gcgctgccgc ccccccgtgg gctgcgagga gctggtgcga gagccgggct  
 240  
 gcggctgttg cgccacttgc gccctgggct tggggatgcc ctgcggggtg tacaccccc  
 300  
 gttgcggctc gggcctgcgc tgctaccgc cccgaggggt ggagaagccc ctgcacacac  
 360  
 tgatgcacgg gcaaggcgtg tgcattggagc tggcggagat cgaggccatc caggaaagcc  
 420  
 tgcagccctc tgacaaggac gagggtgacc accccaacaa cagcttcagc ccctgtagcg  
 480  
 cccatgaccg caggtgcctg cagaagcact tcgccaaaat tcgagaccgg agcaccagtg  
 540  
 ggggcaagat gaaggcctat ggggcgcccc gggaggatgc ccggcctgtg cccaggggct  
 600  
 cctgccagag cgagctgcac cgggcgctgg agcggctggc cgcttcacag agccgcaccc  
 660  
 acgaggacct ctacatcatc cccatcccca actgcgaccg caacggcaac ttccacccca  
 720  
 agcagtgtca cccagctctg gatgggcagc gtggcaagtg ctggtgtgtg gaccggaaga  
 780  
 cgggggtgaa gcttccgggg ggcctggagc caaaggggga gctggactgc caccagctgg  
 840  
 ctgacagctt tcgagagtga ggcctgccag caggccaggg actcagcgtc ccctgtact  
 900  
 cctgtgctct ggaggctgca gagctgacc agagtggagt ctgagtctga gtcctgtctc  
 960  
 tgcctgcggc ccagaagttt ccctcaaagc cgcgtgtgca cgtgtgcgtg tgcgtgcgtg  
 1020  
 tgtgtgtgtt tgtgagcatg ggtgtgccct tggggtaagc cagagcctgg ggtgttctct  
 1080  
 ttggtgttac acagcccaag aggactgaga ctggcactta gcccaagagg tctgagccct  
 1140  
 ggtgtgttcc cagatcgatc ctggattcac tcactcactc attccttcac tcattccagc  
 1200  
 acctaaaaac atttactgac catgtactac gtgccagctc tagttttcag ccttgggagg  
 1260  
 ttttattctg acttctctctg attttggcat gtggagacac tcctataagg agagtccaag  
 1320  
 cctgtgggag tagaaaaatc tcattcccag agtcagagga gaagagacat gtaccttgac  
 1380  
 catcgtcctt cctctcaagc tagccagagg gtgggagcct aaggaagcgt ggggtagcag  
 1440  
 atggagtaat ggtcacgagg tccagacca ctcccaaagc tcagacttgc caggctccct  
 1500  
 ttctcttctt cccaggtcc ttcttttagg tctggttgtt gcaccatctg cttggttggc  
 1560  
 tggcagctga gagccctgct gtgggagagc gaaggggggc aaaggaagac ttgaagcaca  
 1620  
 gagggtcagg gaggtggggt acatttctct gagcagtcag ggtgggaaga aagaatgcaa  
 1680  
 gagggtgactg aatgtgccta atggagaaga cccacgtgct aggggatgag gggcttctctg  
 1740  
 ggtcctgttc cctaccccat ttgtggtcac agccatgaag tcaccgggat gaacctatcc  
 1800  
 ttccagtggc tcgctccctg tagctctgcc tcctctcca tatctccttc cctacacct  
 1860

ccctcccccac acctccctac tcccctgggc atcttctggc ttgactggat ggaaggagac  
 1920  
 ttaggaacct accagttggc catgatgtct ttctt  
 1955

<210> 4  
 <211> 1722  
 <212> DNA  
 <213> Homo sapiens

<400> 4  
 ggggaaaaga gctaggaaag agctgcaaag cagtgtgggc tttttccctt tttttgctcc  
 60  
 ttttcattac ccctcctccg ttttcaccct tctccggact tcgcgtagaa cctgcgaatt  
 120  
 tcgaagagga ggtggcaaag tgggagaaaa gaggtgttag ggtttggggg tttttgtttt  
 180  
 ttgtttttgt tttttaattt cttgatttca acattttctc ccaccctctc ggctgcagcc  
 240  
 aacgcctctt acctgttctg cggcgcccg caccgctggc agctgagggt tagaaagcgg  
 300  
 ggtgtatttt agattttaag caaaaatttt aaagataaat ccatttttct cttccacccc  
 360  
 caacgccatc tccactgcat ccgatctcat tatttcgggtg gttgcttggg ggtgaacaat  
 420  
 tttgtggctt tttttccctt ataattctga cccgctcagg ottgaggggt tctccggcct  
 480  
 ccgctcactg cgtgcacctg gcgctgccct gttccccca acctgttgca aggttttaat  
 540  
 tcttgcaact gggacctgct cgcaggcacc ccagccctcc acctctctct acatttttgc  
 600  
 aagtgtctgg gggagggcac ctgctctacc tgccagaaat tttaaaacaa aaacaaaaac  
 660  
 aaaaaaatct ccggggggccc tcttgggccc tttatccctg cactctcgct ctctgcccc  
 720  
 accccgaggt aaagggggcg actaagagaa gatggtgttg ctcaccgcgg tctcctgct  
 780  
 gctggccgcc tatgcggggc cggcccagag cctgggctcc ttcgtgcaact gcgagccctg  
 840  
 cgacgagaaa gccctctcca tgtgcccccc cagccccctg ggctgcgagc tggtaagga  
 900  
 gccgggctgc ggctgctgca tgacctgcgc cctggccgag gggcagtcgt gcggcgtcta  
 960  
 caccgagcgc tgcgcccagg ggctgcgctg cctcccccg caggacgagg agaagccgct  
 1020  
 gcacgcccctg ctgcacggcc gcgggggttg cctcaacgaa aagagctacc gcgagcaagt  
 1080  
 caagatcgag agagactccc gtgagcacga ggagcccacc acctctgaga tggccgagga  
 1140  
 gacctactcc cccaagatct tccggcccaa acacaccgc atctccgagc tgaaggctga  
 1200  
 agcagtgaag aaggaccgca gaaagaagct gaccagtcct aagtttgtcg ggggagccga  
 1260  
 gaacactgcc ccccccgga tcctctctgc acctgagatg agacaggagt ctgagcaggg  
 1320  
 cccctgcccgc agacacatgg aggttccct gcaggagctc aaagccagcc cacgcatggt  
 1380

gccccgtgct gtgtacctgc ccaattgtga ccgcaaagga ttctacaaga gaaagcagtg  
1440  
caaaccttcc cgtggccgca agcgtggcat ctgctggtgc gtggacaagt acgggatgaa  
1500  
gctgccaggc atggagtacg ttgacgggga ctttcagtgc cacaccttcg acagcagcaa  
1560  
cgttgagtga tgcgtcccc cccaaccttt ccttcacccc ctcccacccc cagccccgac  
1620  
tccagccagc gcctccctcc accccaggac gccactcatt tcatctcatt taagggaaaa  
1680  
atatatatct atctatttga ggaaaaaaaa aaaaaaaaaa aa  
1722



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/08303

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/16 A61K38/18 A61K38/30 A61K38/17 A61K38/39  
 //(A61K38/16, 39:395, 38:17, 31:70)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CARONI P. ET AL: "Signaling by insulin-like growth factors in paralyzed skeletal muscle: Rapid induction of IGF1 expression in muscle fibers and prevention of interstitial cell proliferation by IGF-BP5 and IGF-BP4" THE JOURNAL OF NEUROSCIENCE, vol. 14, no. 5, 1994, pages 3378-3388, XP002115584 cited in the application abstract page 3379, column 1, paragraph 2 page 3385, column 1 page 3386, column 2, paragraphs 2,3 --- -/--	1-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

16 September 1999

Date of mailing of the international search report

05/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Fernandez y Branas, F

# INTERNATIONAL SEARCH REPORT

Internr al Application No

PCT/US 99/08303

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	<p>CIFUENTES-DIAZ C.: "The peripheral nerve and the neuromuscular junction are affected in the tenascin-C deficient mouse"</p> <p>CELLULAR AND MOLECULAR BIOLOGY, vol. 44, no. 2, March 1998 (1998-03), pages 357-379, XP002115585 abstract pages 357-358 "introduction"</p> <p style="text-align: center;">---</p>	12,21
X	<p>BOOTH C.M. ET AL: "An antibody to neural cell adhesion molecule impairs motor nerve terminal sprouting in a mouse muscle locally paralysed with botulinum toxin"</p> <p>NEUROSCIENCE, vol. 35, no. 1, 1990, pages 85-91, XP002115586 abstract page 85, column 1-2</p> <p style="text-align: center;">---</p>	1-8,12, 23
X	<p>GURNEY M.E. ET AL: "Induction of motor neuron sprouting in vivo by ciliary neurotrophic factor and basic fibroblast growth factor"</p> <p>THE JOURNAL OF NEUROSCIENCE, vol. 12, no. 8, 1992, pages 3241-3247, XP002115587 cited in the application the whole document</p> <p style="text-align: center;">---</p>	12,16
X	<p>A. K. MCALLISTER ET AL: "Neurotrophin regulation of cortical dendritic growth requires activity"</p> <p>NEURON, vol. 17, 1996, pages 1057-1064, XP002115588 cited in the application the whole document</p> <p style="text-align: center;">---</p>	12,19
X	<p>CARONI P. ET AL: "Role of muscle insulin-like growth factors in nerve sprouting: suppression of terminal sprouting in paralyzed muscle by IGF-binding protein 4"</p> <p>THE JOURNAL OF CELL BIOLOGY, vol. 125, no. 4, 1994, pages 839-902, XP002115589 cited in the application abstract page 897, paragraph 2 -page 898, paragraph 1</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	1-8, 10-14

# INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/US 99/08303

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 721 215 A (AOKI KEI ROGER ET AL) 24 February 1998 (1998-02-24) column 2, line 20 - line 55 column 5: table 1 column 7; table 1 ---	1-9
T	PAIVA, A. ET AL: "Functional repair of motor endplates after botulinum neurotoxin type A poisoning: Biphasic switch of synaptic activity between nerve sprouts and their parent terminals" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, USA, vol. 96, March 1999 (1999-03), pages 3200-3205, XP002115590 cited in the application the whole document -----	1-24

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/08303

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 1-24 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11

Methods to extend the period during which a tissue treated with a Clostridial toxin is paralysed by contacting the tissue with the compositions mentioned in the said claims 1-11.

2. Claims: 12-24

Methods for stimulating the outgrowth of neural sprouts from damaged neural tissue, comprising contacting the tissue with a composition containing the agents specified in said claims 12-24.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/08303

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5721215 A	24-02-1998	AU 2341797 A	10-10-1997
		CA 2249196 A	25-09-1997
		EP 0889731 A	13-01-1999
		WO 9734624 A	25-09-1997
<hr/>			